

**WEST**[Help](#)[Logout](#)

<a href="#">Main Menu</a>	<a href="#">Search Form</a>	<a href="#">Result Set</a>	<a href="#">Show S Numbers</a>	<a href="#">Edit S Numbers</a>	<a href="#">Referring Patents</a>
<a href="#">First Hit</a>	<a href="#">Previous Document</a>			<a href="#">Next Document</a>	
<a href="#">Full</a>	<a href="#">Title</a>	<a href="#">Citation</a>	<a href="#">Front</a>	<a href="#">Review</a>	<a href="#">Classification</a>
<a href="#">Date</a>	<a href="#">Reference</a>	<a href="#">Claims</a>	<a href="#">KWC</a>		

## Document Number 41

Entry 41 of 53

File: USPT

Jan 3, 1995

DOCUMENT-IDENTIFIER: US 5378464 A

TITLE: Modulation of inflammatory responses by  
administration of GMP-140 or antibody to GMP-140

## DEPR:

Other studies have demonstrated that GMP-140 binds to C3b, a complement system protein, and protein S, an anticoagulant cofactor protein. GMP-140 shares sequence homology with the plasma protein C4b-binding protein (C4bp), which not only interacts with the plasma protein C4b but also with protein S.

## DEPR:

One of the complement-regulatory proteins containing repeating units homologous to those in GMP-140 is C4b-binding protein (C4bp). In addition to binding C4b, this plasma protein also binds the anticoagulant cofactor, protein S. Protein S serves as a cofactor for the anticoagulant serine protease, activated protein C. Activated protein C proteolytically inactivates coagulation Factors V and VIII, thus suppressing the coagulation cascade. Binding of protein S to C4bp blocks the anticoagulant functions of protein S. The dual binding specifications of C4bp for C4b and protein S suggested that GMP-140 might also bind protein S. This possibility was especially attractive because certain vascular cell receptors such as Mac-1 on monocytes can have several different functions related to both inflammation and coagulation. A number of studies have now shown that purified GMP-140 does interact with purified protein S. Initial studies suggest that the binding facilitates the anticoagulant cofactor function of protein S. Accordingly, infusion of soluble fragments of GMP-140 containing the protein S-binding site might serve as novel anticoagulants.

<a href="#">Main Menu</a>	<a href="#">Search Form</a>	<a href="#">Result Set</a>	<a href="#">Show S Numbers</a>	<a href="#">Edit S Numbers</a>	<a href="#">Referring Patents</a>
<a href="#">First Hit</a>	<a href="#">Previous Document</a>			<a href="#">Next Document</a>	
<a href="#">Full</a>	<a href="#">Title</a>	<a href="#">Citation</a>	<a href="#">Front</a>	<a href="#">Review</a>	<a href="#">Classification</a>
<a href="#">Date</a>	<a href="#">Reference</a>	<a href="#">Claims</a>	<a href="#">KWC</a>		

[Help](#)[Logout](#)

**WEST**[Help](#)[Logout](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show 8 Numbers](#)[Edit 8 Numbers](#)**Search Results -**

Term	Documents
CD4.USPT.	3294
CD4S.USPT.	4
(1 AND CD4).USPT.	18

Database: 

Refine Search:

**Search History**

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT	11 and cd4	18	<u>L7</u>
USPT	14 and cd4	10	<u>L6</u>
USPT	14 and (chimeric or chimreras or multimeric or multimer)	16	<u>L5</u>
USPT	c4bp	53	<u>L4</u>
USPT	11 and (multimeric or multimer)	10	<u>L3</u>
USPT	11 (multimeric or multimer)	0	<u>L2</u>
USPT	c4bp or proline-rich protein or proline rich protein	101	<u>L1</u>